

# budesonide + formoterol

## New dose regimen

### Asthma attacks: a step backwards

● **Using the budesonide + formoterol combination for both maintenance treatment of asthma and on-demand control of asthma attacks amounts to intensifying maintenance treatment. It is not a valid treatment objective, and is likely to confuse asthma patients.**



NOT ACCEPTABLE

Maintenance and on-demand treatment of asthma are based on the results of many clinical trials involving several hundred thousand patients (1-3). Maintenance treatment depends on asthma severity and is based on inhaled steroid monotherapy at the minimal effective dose. Adding an inhaled long-acting beta-2 agonist does not seem to provide a tangible clinical benefit for patients (1,2).

Treatment of asthma attacks is based on short-acting high-dose beta-2 agonists, which are rapidly effective and have few adverse effects (4,5).

The fixed-dose combination of budesonide and formoterol (Symbicort<sup>o</sup>, AstraZeneca) was first marketed for maintenance treatment of asthma. The SPC now also mentions, in the posology section, the possible use of this product to treat asthma attacks.

**A two-phase licence extension process.** The company first submitted, through the European mutual recognition procedure, an application including data from three trials. These trials compared budesonide + formoterol, for both maintenance and on-demand treatment, versus budesonide, alone or with formoterol, for maintenance treatment, and terbutaline for attacks (6-10). The application was rejected in 2004 (9). The company then submitted new data, and the new marketing terms were finally granted in late 2006 (10).

**Intensified maintenance treatment.** The new data are mainly based on two double-blind randomised trials including 3394 and 3335 patients (11,12). The first trial, in patients who were all receiving maintenance treatment with budesonide + formoterol, compared treatment of asthma attacks with budesonide + formoterol, or formoterol, or terbutaline (11). The second trial also included

three groups: two groups received maintenance treatment with budesonide + formoterol, and used budesonide + formoterol or terbutaline for asthma attacks; the third group received maintenance treatment with fluticasone + salmeterol and used terbutaline for attacks (12).

In these trials the patients used their "attack" treatments once or twice a day on average, which, in the groups only treated with budesonide + formoterol, represented an intensification of the maintenance treatment. It is therefore hardly surprising that patients in these groups showed more improvement in some criteria for asthma control (11,12).

#### Increased risk of adverse effects.

There were no major differences in adverse effects between the different treatments tested in these trials (6-8,11,12). However, the use of the budesonide + formoterol combination for both maintenance and on-demand treatment leads to an increase in drug exposure, and a higher risk of adverse effects (a).

Long-acting beta-2 agonists accumulation may carry an added risk of cardiac disorders, tremor, hypokalaemia and hyperglycaemia (13).

Unless combined with an inhaled steroid, long-acting beta-2 agonists can increase the risk of hospitalisation and life-threatening asthma attacks. There are reports of asthma aggravation associated with the use of a long-acting beta-2 agonist (salmeterol), even when combined with a steroid (2,14).

In addition, the risk of adrenal failure increases with the dose of inhaled steroid (including budesonide) (15).

The budesonide + formoterol combination should not be recommended for 'on demand' use to control asthma attacks (b). It exposes patients to an added risk of adverse effects, and undermines the clear-cut message that has been found to help patients manage their asthma: the clear distinction between maintenance treatment, with an inhaled steroid at the minimal required dose, and control of attacks with a short-acting beta-2 agonist.

©Prescrire

a- A risk management plan has been launched (ref 16).  
b- The French Transparency Committee concluded that,

### budesonide + formoterol (Symbicort<sup>o</sup>)

Powder for inhalation

■ **New posology:** "Adults (18 years and older) (...): Patients should take 1 additional inhalation as needed in response to symptoms. If symptoms persist after a few minutes, an additional inhalation should be taken. (...) a total daily dose of up to 12 inhalations could be used for a limited period (...)"

[French marketing authorisation through the mutual recognition procedure after initial Swedish authorisation]

#### Steroid + long-acting beta-2 agonist

in this setting, Symbicort<sup>o</sup> did not represent an improvement over existing treatments (ref 17).

#### Selected references from *Prescrire's* literature search.

- 1- Prescrire Rédaction "Traitement de l'asthme - 1<sup>re</sup> partie" *Rev Prescrire* 2007; 27 (284): 436-441.
- 2- Prescrire Editorial Staff "Long-term asthma therapy" *Prescrire Int* 2007; 16 (91): 522-526.
- 3- British Thoracic Society and Scottish Intercollegiate Guidelines Network "British Guideline on the management of asthma. A national Clinical Guideline" Revised ed., November 2005: 94 pages.
- 4- Prescrire Editorial Staff "How to treat asthma attacks" *Prescrire Int* 1996; 5 (25): 154-158.
- 5- Prescrire Rédaction "Crise d'asthme grave à domicile" *Rev Prescrire* 2007; 27 (284): 441-444.
- 6- Rabe KF et al. "budesonide/formoterol in a single inhaler for maintenance and relief in mild-to-moderate asthma: a randomized, double blind trial" *Chest* 2006; 129: 246-256.
- 7- Scicchitano R et al. "Efficacy and safety of budesonide/formoterol single inhaler therapy versus a higher dose of budesonide in moderate to severe asthma" *Curr Med Res Opin* 2004; 20 (9): 1403-1418.
- 8- Byrne P "budesonide/formoterol combination therapy as both maintenance and reliever medication in asthma" *Am J Respir Crit Care Med* 2005; 171: 129-136.
- 9- "AstraZeneca hit by EU delay for Symbicort single inhaler therapy" *Scrip* 2004; (3002): 22.
- 10- "Symbicort Smart finally gets the nod in Europe" *Scrip* 2006; (3200): 19.
- 11- Rabe KF et al. "Effect of budesonide in combination with formoterol for reliever therapy in asthma exacerbations: a randomised controlled, double-blind study" *Lancet* 2006; 368: 744-753.
- 12- Kuna P et al. "Effect of budesonide/formoterol maintenance and reliever therapy on asthma exacerbations" *Int J Clin Pract* 2007; 61 (5): 725-736.
- 13- Prescrire Rédaction "18-1-2. Patients sous bêta-2 stimulant" *Rev Prescrire* 2007; 27 (290 suppl. interactions médicamenteuses).
- 14- Prescrire Editorial Staff "Asthma: fatal exacerbation with long-acting beta 2 agonist therapy without concomitant steroid therapy" *Prescrire Int* 2007; 16 (88): 69-70.
- 15- Prescrire Rédaction "Insuffisance surrénale aiguë: avec les corticoïdes inhalés aussi" *Rev Prescrire* 2007; 27 (289): 826.
- 16- AFSSAPS "PGR-Symbicort" July 2007: 2 pages.
- 17- HAS - Commission de la transparence "Avis-Symbicort" 4 July 2007: 14 pages.